



How to manage your winemaking process with grapes affected by fire?

Treatment protocol during winemaking

ON GRAPE

Harvest

Favor manual harvesting, if mechanical harvesting:

use a double bottom trailer, separate the draining juice from the grapes and get rid of it.

WHITE AND ROSÉ WINEMAKING PROCESS

DIRECT PRESSING

Clarify by depectinization as usual:
Viazym® Clarif Extrem at 0.5 mL/hL

STEP 1

Use **Provgreen® Smoke Taints** 40 g/hL wait 12 hours then rack

STEP 2

Yeast + nutrition (as usual)
AT MID FERMENTATION (D 1020)
Add **Viazym® Aroma** 3 g/hL

STEP 3

To prevent inhibiting the enzyme wait 10 days before adding bentonite, then rack according to bentonite recommendations

STEP 4

Treat with **Provgreen® Smoke Taints** 40 g/hL, wait 24 hours then rack

STEP 5

Taste! If smoke taints are still presents, repeat step 4 with 20 g/hL of **Provgreen® Smoke Taints**

RED CLASSICAL WINEMAKING PROCESS

STEP 1

Vatting of the harvest, during the pumping over for the tank homogenization add **Provgreen® Smoke Taints** 40 g/hL. No racking **Provgreen® Smoke Taints** can remain throughout the alcoholic fermentation

STEP 2

Yeast + nutrition (as usual). Pump over the tank 2 times per day for 3 days - then the following days pump over according to cellar protocol, favor short maceration max 5 days

PRESSING

STEP 3

Finish AF in liquid phase and add **Viazym® Aroma** at 3 g/hL After AF finished wait 10 days then rack

STEP 4

Add **Provgreen® Smoke Taints** 40 g/hL, wait 24 hours then rack

STEP 5

Taste! If smoke taints are still presents, repeat step 4 with 20 g/hL of **Provgreen® Smoke Taints**

Maximum legal dose for use:
100 g/hL

Provgreen® Smoke Taints:

Preparation to be mix exclusively with water as 1 kg per 10 L of water!



IMPORTANT

Take care to homogenize the tank well after each addition of **Provgreen® Smoke Taints**, minimum pump over twice the total volume of the tank

DOSAGE

Recommended dosage:

40 to 100 g/hl depending on contamination level.

Maximum legal dose according to current European regulations: 100 g/hl.

Pre-testing is necessary to determine the optimal dose.

INSTRUCTIONS FOR USE

Disperse the product in 10 times its weight in water (do not use must). Homogenize. Then add to the tank when pumping over (the use of a fining connector is recommended).

Rack the wine 48 to 72 hours after treatment.

Please refer to dedicated protocols for treatment

Precautions for use:

For oenological and exclusively professional use.

Use in accordance with current regulations.

PACKAGING

5 kg Bag

STORAGE

Unused, sealed original package: keep in a dry, odourless place away from light.

Once opened, use up within 48 hours.

The information above corresponds to the current state of our knowledge. It does not represent a commitment or guarantee, since the conditions of use are beyond our control. It does not exempt the user from responsibility in complying with the legislation and safety standards in force. This document is the property of SOFRALAB and may not be amended without its consent.